

# Prevantics® Maxi Swabstick

Skin Antisepsis

3.15% Chlorhexidine Gluconate (w/v) and 70% Isopropyl Alcohol (v/v)  
Preinjection & Preoperative Skin Preparation

## INSTRUCTIONS FOR USE

- For Item No. S41950: Peel apart the pouch to expose the swabstick.
- For Item No. S27350 (kit): Grasp the pouch at the notch and tear open straight across.
- Remove the swabstick from package. Do not touch the foam tip.



**PEEL APART**  
Item No. S41950



**TEAR OPEN**  
Item No. S27350 (kit)

## APPLY

- For dry sites: Scrub repeatedly back and forth for 15 seconds with each side of the applicator for a total of 30 seconds. Allow to air dry for approximately 30 seconds.
- For moist sites: Scrub repeatedly back and forth for 1 minute with each side of the applicator for a total of 2 minutes. Allow to air dry for approximately 1.5 minutes.
- Maximum treatment area for one swabstick:
  - **Dry Site: 7 x 7 in. (18 x 18 cm.)**
  - **Moist Site: 3 x 7.5 in. (7.5 x 19 cm.)**
- Do not blot or wipe dry.
- Discard after single use.



See Drug Facts panel for complete product information.

**REORDER #** \_\_\_\_\_ **REPLACING ORDER #** \_\_\_\_\_

1-800-999-6423 + Phone: (845) 365-1700 + Fax: (845) 398-5347 + www.PDIHC.com

©2018 PDI.  
PDI02189756

**PDI**  
BE THE DIFFERENCE